

**Texas Allergy** 

Phone: (281) 886-7440 Fax: (281) 929-0086 info@texasallergygroup.com

http://texasallergygroup.com

# MEDICATIONS THAT NEED TO BE STOPPED FOR ALLERGY TESTING

THE FOLLOWING IS A LIST OF MEDICATIONS THAT YOU **SHOULD STOP TAKING** PRIOR TO ALLERGY TESTING. THIS ALSO INCLUDES THE LENGTH OF TIME TO AVOID THESE MEDICATIONS.

## STOP THE FOLLOWING MEDICATIONS ONE WEEK PRIOR TO TESTING:

CLARITIN, CLARITIN-D, CLARINEX
LORATADINE (GENERIC FOR CLARITIN)
ZYRTEC (CETIRIZINE), ZYRTEC-D, XYZAL (LEVOCETIRIZINE)
ALLEGRA AND ALLEGRA-D
FEXOFENADINE (GENERIC TO ALLEGRA)
ATARAX OR HYDROXYZINE
CYPROHEPTADINE OR PERIACTIN
TAVIST (CLEMASTINE)
ASTELIN NASAL SPRAY
OPTIVAR

AND ALL OTHER UNKNOWN BRAND OF OVER THE COUNTER COUGH/COLD/FLU/SINUS MEDICATIONS

## STOP THE FOLLOWING MEDICATIONS AT LEAST 3 DAYS PRIOR TO TESTING:

ZANTAC or RANITIDINE PEPCID OR FAMOTIDINE AXID OR NIZATIDINE TAGAMET OR CIMETIDINE

BENADRYL OR DIPHENHYDRAMINE CHLORPHENIRAMINE OR CHLOR-TRIMETON DIMETAPP PHENERGAN TYLENOL PM UNISOM

# STOP THE FOLLOWING MEDICATIONS AT MIDNIGHT THE DAY PRIOR TO YOUR APPOINTMENT:

ACCOLATE (ZAFIRLUKAST) REGLAN (METOCLOPRAMIDE) SINGULAIR (MONTELUKAST) ZYFLO (ZILEUTIN)

There are other drugs that need to be stopped for allergy testing but require your prescribing physician's approval before stopping. Your medication list will be reviewed by your allergist at your first appointment and you will be notified if you are taking any of these medications. Arrangements can then be made with the prescriber to see if you can stop the medication for allergy testing. Also, you do not need to stop antihistamines if you have chronic hives. Please e-mail <a href="info@texasallergygroup.com">info@texasallergygroup.com</a> or call clinic at 281-886-7440 if you have any questions about which medications to stop prior to your visit.

## THE FOLLOWING MEDICATIONS ARE OKAY TO USE PRIOR TO TESTING:

ALL INHALERS FOR ASTHMA, NASAL STEROID SPRAYS AND PREDNISONE