



Texas Allergy

Clinic for Allergy Asthma Sinus and Immunologic Diseases

Phone: (281) 886-7440

Fax: (281) 929-0086

E-mail: info@texasallergygroup.com

<http://www.texasallergygroup.com>

MEDICATIONS THAT NEED TO BE STOPPED FOR ALLERGY TESTING

THE FOLLOWING IS A LIST OF MEDICATIONS THAT YOU ***SHOULD STOP TAKING*** PRIOR TO ALLERGY TESTING. THIS ALSO INCLUDES THE LENGTH OF TIME TO AVOID THESE MEDICATIONS.

STOP THE FOLLOWING MEDICATIONS ONE WEEK PRIOR TO TESTING:

CLARITIN, CLARITIN-D, CLARINEX
LORATADINE (GENERIC FOR CLARITIN)
ZYRTEC (CETIRIZINE), ZYRTEC-D, XYZAL (LEVOCETIRIZINE)
ALLEGRA AND ALLEGRA-D
FEXOFENADINE (GENERIC TO ALLEGRA)
ATARAX OR HYDROXYZINE
CYPROHEPTADINE OR PERIACTIN
TAVIST (CLEMASTINE)
ASTELIN NASAL SPRAY, OPTIVAR

AND ALL OTHER UNKNOWN BRAND OF OVER THE COUNTER COUGH/COLD/FLU/SINUS MEDICATIONS

STOP THE FOLLOWING MEDICATIONS AT LEAST 3 DAYS PRIOR TO TESTING:

ZANTAC or RANITIDINE
PEPCID OR FAMOTIDINE
AXID OR NIZATIDINE
TAGAMET OR CIMETIDINE

BENADRYL OR DIPHENHYDRAMINE
CHLORPHENIRAMINE OR CHLOR-TRIMETON
DIMETAPP
PHENERGAN
TYLENOL PM
UNISOM

STOP THE FOLLOWING MEDICATIONS AT MIDNIGHT THE DAY PRIOR TO YOUR APPOINTMENT:

ACCOLATE (ZAFIRLUKAST)
REGLAN (METOCLOPRAMIDE)
SINGULAIR (MONTELUKAST)
ZYFLO (ZILEUTIN)

There are other drugs that need to be stopped for allergy testing but require your prescribing physician's approval before stopping. Your medication list will be reviewed by your allergist at your first appointment and you will be notified if you are taking any of these medications. Arrangements can then be made with the prescriber to see if you can stop the medication for allergy testing. Also, you do not need to stop antihistamines if you have chronic hives. Please e-mail info@texasallergygroup.com or call clinic at 281-886-7440 if you have any questions about which medications to stop prior to your visit.

THE FOLLOWING MEDICATIONS ARE OKAY TO USE PRIOR TO TESTING:

ALL INHALERS FOR ASTHMA, NASAL STEROID SPRAYS AND PREDNISONE